

ON THE MODULI PART OF THE KAWAMATA-KODAIRA CANONICAL BUNDLE FORMULA

CAUCHER BIRKAR

YIFEI CHEN

ABSTRACT. It is conjectured that the moduli b-divisor of the Kawamata-Kodaira canonical bundle formula associated to a klt-trivial fibration $(X, B) \rightarrow Z$ is semi-ample. In this paper, we show the semi-ampleness of an arbitrarily small perturbation of the moduli b-divisor by a fixed appropriate divisor which roughly speaking comes from a section of $K_X + B$.

We apply the above result to settle a conjecture of Fujino and Gongyo: if $f: X \rightarrow Z$ is a smooth surjective morphism of smooth projective varieties with $-K_X$ semi-ample, then $-K_Z$ is also semi-ample. We list several counter-examples to show that this fails without the smoothness assumption on f .

1. INTRODUCTION

The semi-ampleness conjecture. We work over $\mathbb{C}$. Let $f: X \rightarrow Z$ be a contraction of normal projective varieties, and (X, B) klt such that $K_X + B \sim_{\mathbb{Q}} 0/Z$, i.e. $K_X + B \sim_{\mathbb{Q}} f^*N$ for some $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor N . Such a contraction is called a *klt-trivial fibration*. By a construction of Kawamata [Ka97],[Ka98] we have a decomposition

$$N \sim_{\mathbb{Q}} K_Z + B_Z + M_Z$$

where B_Z is defined using the singularities of (X, B) and of the fibres of f over the codimension one points of Z , and (Z, B_Z) is klt if $K_Z + B_Z$ is $\mathbb{Q}$ -Cartier. The part B_Z is called the *discriminant part* and the part M_Z is called the *moduli part*. More precisely, B_Z is defined as follows: for each prime divisor Q on Z , let t be the lc threshold of f^*Q over the generic point of Q , with respect to the pair (X, B) ; then let $(1 - t)$ be the coefficient of Q in B_Z . Except for finitely many Q , $t = 1$ hence B_Z has finitely many components.

Consider a commutative diagram

$$\begin{array}{ccc} X' & \xrightarrow{\tau} & X \\ \downarrow f' & & \downarrow f \\ Y & \xrightarrow{\sigma} & Z \end{array}$$

in which X', Y are normal and projective, σ, τ are birational, and f' is a contraction. Let $K_{X'} + B' := \tau^*(K_X + B)$ where B' might have negative coefficients. Using the relation $K_{X'} + B' \sim_{\mathbb{Q}} 0/Y$, we can similarly define a decomposition

$$\sigma^*N \sim_{\mathbb{Q}} K_Y + B_Y + M_Y$$

Date: July 18, 2012.

where: $B_Z = \sigma_* B_Y$ and $M_Z = \sigma_* M_Y$. This means that the moduli part is a b-divisor. In general, B_Y may have negative coefficients.

Kawamata [Ka98] showed that M_Y is nef if Y is a sufficiently high resolution. Ambro [Am04] proved that the moduli part is a b-Cartier divisor, i.e. we can find a resolution $Y \rightarrow Z$ so that for any other resolution $Y' \rightarrow Y$ the moduli part $M_{Y'}$ is just the pullback of M_Y . Moreover, Ambro [Am05] proved that M_Y is actually the pullback of some nef and big divisor for some contraction $g: Y \rightarrow T$. We call Y an *Ambro model*.

Conjecture 1.1 (cf. [Ka97], [Am99], [PSh09]). *If Y is an Ambro model, then M_Y is semi-ample.*

This seems to be a difficult conjecture. Only some very special cases are known: when $M_Y \equiv 0$ [Am05], when $\dim X = \dim Z + 1$ [Ka97][PSh09], and when the geometric generic fibre of f is birationally an abelian variety [F03].

Results of this paper. We are not able to prove the semi-ampleness of M_Y but we prove semi-ampleness of a "nearby" divisor. This would be enough for some applications. To state our main result concerning M_Y we need to introduce one more notation. Since (Y, B_Y) is log smooth and each coefficient of B_Y is strictly smaller than 1, there is an exceptional/ Z $\mathbb{Q}$ -divisor $E \geq 0$ on Y such that (Y, Δ_Y) is klt where $\Delta_Y := B_Y + E \geq 0$.

Theorem 1.2. *Under the above notation and assumptions, suppose that we have $\kappa(K_Y + \Delta_Y/T) \geq 0$, i.e. suppose that there is a rational number $b \geq 0$ such that*

$$K_Y + \Delta_Y + bM_Y \sim_{\mathbb{Q}} D_Y \text{ for some } D_Y \geq 0$$

Then, we can choose $D_Y \geq 0$ in its $\mathbb{Q}$ -linear equivalence class so that there exist a resolution $\phi: W \rightarrow Y$ and a $\mathbb{Q}$ -divisor G on W such that

- $0 \leq G \leq \phi^* D_Y$,
- $\alpha M_W + G = \alpha \phi^* M_Y + G$ is semi-ample for any $\alpha \gg 0$, and
- $R(W, lG) \simeq R(Y, lD_Y)$ for some integer $l > 0$.

Here $R(-)$ stands for the section ring (see Section 2 for precise definition). The proof is based on the minimal model program. An interesting special case of the theorem is when $K_X + B \sim_{\mathbb{Q}} 0$ in which case we take $D_Y = E$. Even this special case is strong enough to imply the following result which was conjectured by Fujino and Gongyo [FG11] (see also [FG12]).

Theorem 1.3. *Let $f: X \rightarrow Z$ be a smooth surjective morphism of smooth projective varieties. If $-K_X$ is semi-ample, then $-K_Z$ is also semi-ample.*

Fujino-Gongyo proved the conjecture when $-K_X$ is big. They also proved some other special cases (see Remark 4.2 and Theorem 4.4 of [FG11]). The theorem does not hold without the smoothness assumption on f by a counter-example due to H. Sato (see Section 4 where we give additional counter-examples).

Relevant results. There are results similar to Theorem 1.3 in the literature. We mention some of them for completeness. Let $f: X \rightarrow Z$ be a smooth

surjective morphism of smooth projective varieties. Then, it is known that (see [FG11], §5 for more details):

- if $-K_X$ is nef, then $-K_Z$ is nef (Miyaoka [M93])
- if $-K_X$ is ample, then $-K_Z$ is ample (Kollár-Miyaoka-Mori [KMM92])
- if $-K_X$ is nef and big, then $-K_Z$ is nef and big (Fujino-Gongyo [FG11]).

To prove Theorem 1.3, we do not rely on these known results but we use ideas of [FG11].

In general, varieties with nef anticanonical divisor have interesting properties. Let X be a smooth projective variety with $-K_X$ nef. Then, for any surjective morphism $g: X \rightarrow Y$ with Y smooth projective, Zhang [Zh96] proved that either Y is uniruled or $\kappa(Y) = 0$. On the other hand, Lu, Tu, Zhang, and Zheng [LTZZ10] proved that the Albanese map $X \rightarrow A$ is semi-stable (it is conjectured that this map should be smooth).

Acknowledgements. We would like to thank Osamu Fujino and Florin Ambro for their comments and answering our questions. This work started when Caucher was visiting Yifei at the Mathematics Institute of the Chinese Academy of Sciences in Beijing; he would like to thank them for their hospitality. This work was completed when Yifei was visiting Caucher in Cambridge and he would like to thank Caucher and his family for the hospitality. Both authors were supported by a Leverhulme grant.

2. PRELIMINARIES

Pairs. We work over $k = \mathbb{C}$. A *pair* (X, B) consists of a normal quasi-projective variety X over k and a $\mathbb{Q}$ -divisor B on X with coefficients in $[0, 1]$ (called a boundary) such that $K_X + B$ is $\mathbb{Q}$ -Cartier. A pair (X, B) is called *Kawamata log terminal* (klt) if for any projective birational morphism $g: Y \rightarrow X$ from a normal variety Y , every coefficient of B_Y is less than one where $K_Y + B_Y := g^*(K_X + B)$. For basic properties of singularities and other aspects of birational geometry we refer the reader to [KM98].

Minimal models. Let X be a normal projective variety and D a $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor on X . A normal projective variety Y is called a *minimal model* of D if there is a birational map $\phi: X \dashrightarrow Y$ such that ϕ^{-1} does not contract divisors, Y is $\mathbb{Q}$ -factorial, $D_Y := \phi_* D$ is nef, and there is a common resolution $g: W \rightarrow X$ and $h: W \rightarrow Y$ such that $E := g^* D - h^* D_Y$ is effective and $\text{Supp } g_* E$ is equal to the union of the exceptional divisors of ϕ . If D_Y is semi-ample, we call Y a *good* minimal model of D . A standard way of obtaining a minimal model of D is to run a *minimal model program* on D , if possible. Such a program can be defined exactly as in the case $D = K_X + B$ for a klt pair (X, B) .

When (X, B) is klt and $D = K_X + B$ the above definition of minimal model is equivalent to the usual definition of log minimal models. Moreover, if $K_X + B$ is pseudo-effective and B is big, then $K_X + B$ has a good log minimal model by [BCHM].

Section rings. For a $\mathbb{Q}$ -divisor D on a normal projective variety X , the *section ring* of D is defined as

$$R(X, D) = \bigoplus_{0 \leq m \in \mathbb{Z}} H^0(X, \lfloor mD \rfloor)$$

If $\phi: X \dashrightarrow Y$ is a partial minimal model program on D , i.e. a finite sequence of divisorial contractions and flips with respect to D , then there is a common resolution $g: W \rightarrow X$ and $h: W \rightarrow Y$ such that $E := g^*D - h^*D_Y$ is effective and $\text{Supp } g_*E$ is equal to the union of the exceptional divisors of ϕ . In particular, this implies that there is a natural isomorphism

$$R(X, D) \simeq R(Y, D_Y)$$

Kodaira dimension. Let $f: X \rightarrow Z$ be a contraction of normal varieties and D a $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor on X . By $\kappa(D/Z)$ we mean the Kodaira dimension of $D|_F$ where F is the generic fibre of f .

3. PROOF OF THEOREM 1.2

In this section, we prove the following result which is of independent interest and it immediately implies Theorem 1.2. The proof also works for $\mathbb{R}$ -divisors but for simplicity we only consider $\mathbb{Q}$ -divisors.

Proposition 3.1. *Let (Y, Δ_Y) be a projective klt pair with Δ_Y a $\mathbb{Q}$ -divisor, $g: Y \rightarrow T$ a contraction, M_T a nef and big $\mathbb{Q}$ -divisor on T , and $M_Y = g^*M_T$. Assume that $b \geq 0$ is a rational number such that*

$$K_Y + \Delta_Y + bM_Y \sim_{\mathbb{Q}} D_Y \text{ for some } D_Y \geq 0$$

Then, we can choose $D_Y \geq 0$ in its $\mathbb{Q}$ -linear equivalence class such that there exist a resolution $\phi: W \rightarrow Y$ and a $\mathbb{Q}$ -divisor G on W such that

- $0 \leq G \leq \phi^*D_Y$,
- $\alpha\phi^*M_Y + G$ is semi-ample for any $\alpha \gg 0$, and
- $R(W, lG) \simeq R(Y, lD_Y)$ for some integer $l > 0$.

Proof. For any morphism $U \rightarrow T$, M_U will denote the pullback of M_T .

Step 1. Let $\pi: Y' \rightarrow Y$ be any log resolution of (Y, Δ_Y) . There is an exceptional/ Y $\mathbb{Q}$ -divisor $E' \geq 0$ and a boundary $\Delta_{Y'}$ such that $(Y', \Delta_{Y'})$ is klt and

$$K_{Y'} + \Delta_{Y'} = \pi^*(K_Y + \Delta_Y) + E'$$

Let $D_{Y'} = \pi^*D_Y + E'$. Then,

$$K_{Y'} + \Delta_{Y'} + bM_{Y'} \sim_{\mathbb{Q}} D_{Y'}$$

Now assume that the result holds on Y' , that is, assume that we can choose $D_{Y'} \geq 0$ in its $\mathbb{Q}$ -linear equivalence class such that there exist a resolution $\psi: W \rightarrow Y'$ and a $\mathbb{Q}$ -divisor G on W satisfying:

- $0 \leq G \leq \psi^*D_{Y'}$,
- $\alpha M_W + G$ is semi-ample for any $\alpha \gg 0$, and

- $R(W, lG) \simeq R(Y', lD_{Y'})$ for some integer $l > 0$.

We will show that the result also holds for Y by taking $\phi = \pi\psi$. Since $D_{Y'} = \pi^*D_Y + E'$ and $E' \geq 0$ is exceptional/ Y ,

$$R(Y, lD_Y) \simeq R(Y', lD_{Y'}) \simeq R(W, lG)$$

It only remains to show that $G \leq \phi^*D_Y$. Put $C = \phi^*D_Y - G$. Since E' is exceptional/ Y ,

$$\phi_*C = \phi_*(\phi^*D_Y - G) = \phi_*(\phi^*D_Y + \psi^*E' - G) = \phi_*(\psi^*D_{Y'} - G) \geq 0$$

On the other hand, $\alpha M_W + G$ is semi-ample for any $\alpha \gg 0$ and $M_W \sim_{\mathbb{Q}} 0/Y$ hence G is semi-ample over Y . So, C is antinef over Y which implies that $C \geq 0$ by the negativity lemma. Therefore, $G \leq \phi^*D_Y$.

Step 2. Since $M_Y \sim_{\mathbb{Q}} 0/T$, $K_Y + \Delta_Y \sim_{\mathbb{Q}} D_Y/T$ hence $\kappa(K_Y + \Delta_Y/T) \geq 0$. By taking a log resolution and applying Step 1 we could assume that the relative Iitaka fibration of $K_Y + \Delta_Y$ over T is a morphism $Y \rightarrow S'/T$ where S' is smooth. In particular, this means that $\kappa(K_Y + \Delta_Y/S') = 0$. By applying Fujino-Mori [FM00], we can find a commutative diagram

$$\begin{array}{ccc} & V & \\ \tau \swarrow & & \searrow h \\ Y & & S \\ g \searrow & & \swarrow \theta \\ & T & \end{array}$$

where τ is a resolution, and h, θ are contractions such that we have the following data:

- a klt (S, Δ_S) where Δ_S is a $\mathbb{Q}$ -divisor,
- a nef $\mathbb{Q}$ -divisor L_S on S ,
- $K_S + \Delta_S + L_S$ is big/ T ,
- $\kappa(K_Y + \Delta_Y/T) = \kappa(K_S + \Delta_S + L_S/T) = \dim S - \dim T$,
- and

$$\tau^*(K_Y + \Delta_Y) + R^- \sim_{\mathbb{Q}} h^*(K_S + \Delta_S + L_S) + R^+$$

where R^+, R^- are effective $\mathbb{Q}$ -divisors with R^- exceptional/ Y , $h(R^-)$ has no codimension one components, and $h_*\mathcal{O}_V([iR^+]) = \mathcal{O}_S$ for every $i > 0$.

The above properties imply that for any rational number $\alpha \geq 0$, we have

$$\begin{aligned} \tau^*D_Y + R^- + \alpha M_V &\sim_{\mathbb{Q}} \tau^*(K_Y + \Delta_Y + bM_Y) + R^- + \alpha M_V \\ &= \tau^*(K_Y + \Delta_Y) + R^- + bM_V + \alpha M_V \\ &\sim_{\mathbb{Q}} h^*(K_S + \Delta_S + L_S) + R^+ + bM_V + \alpha M_V \\ &= h^*(K_S + \Delta_S + L_S + bM_S + \alpha M_S) + R^+ \end{aligned}$$

Step 3. Fix a rational number $\lambda \geq 0$ so that $b + \lambda > 0$. By construction

$$0 \leq \kappa(D_Y) = \kappa(K_Y + \Delta_Y + bM_Y) = \kappa(K_S + \Delta_S + L_S + bM_S)$$

so there is a $\mathbb{Q}$ -divisor $D_S \geq 0$ such that

$$D_S \sim_{\mathbb{Q}} K_S + \Delta_S + L_S + bM_S$$

Since D_S is big/ T and M_T is big, there is $a > 0$ such that $D_S + aM_S$ is big. Thus, we have

$$D_S + aM_S \sim_{\mathbb{Q}} A + B$$

where A is ample and $B \geq 0$.

Pick $\epsilon > 0$ sufficiently small. This ensures that

$$\beta_\lambda := \epsilon\lambda - \epsilon a + b + \lambda \geq 0$$

Moreover, since A is ample and L_S and M_S are nef, there is a big boundary $\Gamma_{S,\lambda}$ such that $(S, \Gamma_{S,\lambda})$ is klt and such that we can write

$$\begin{aligned} K_S + \Gamma_{S,\lambda} &\sim_{\mathbb{Q}} K_S + \Delta_S + L_S + \epsilon A + \epsilon B + \beta_\lambda M_S \\ &\sim_{\mathbb{Q}} K_S + \Delta_S + L_S + \epsilon D_S + \epsilon a M_S + \beta_\lambda M_S \\ &\sim_{\mathbb{Q}} K_S + \Delta_S + L_S + b M_S + \epsilon D_S + \epsilon \lambda M_S + \lambda M_S \\ &\sim_{\mathbb{Q}} (1 + \epsilon)(K_S + \Delta_S + L_S + b M_S) + (1 + \epsilon)\lambda M_S \\ &\sim_{\mathbb{Q}} (1 + \epsilon)(K_S + \Delta_S + L_S + b M_S + \lambda M_S) \\ &\sim_{\mathbb{Q}} (1 + \epsilon)(D_S + \lambda M_S) \end{aligned}$$

By [BCHM], we can run an LMMP on $K_S + \Gamma_{S,\lambda}$ which ends up with a good log minimal model of $K_S + \Gamma_{S,\lambda}$. By the relations above, this also produces a good minimal model of $K_S + \Delta_S + L_S + b M_S + \lambda M_S$.

For any $\alpha \geq \lambda$, we have


$$K_S + \Gamma_{S,\alpha} \sim_{\mathbb{Q}} K_S + \Gamma_{S,\lambda} + (1 + \epsilon)(\alpha - \lambda)M_S$$

If $\alpha \gg \lambda$ and if we run an LMMP on $K_S + \Gamma_{S,\alpha}$, then M_S is numerically trivial on each extremal ray contracted in the process: this follows from the boundedness of the length of extremal rays due to Kawamata [Ka91]; indeed if R is an extremal ray such that $(K_S + \Gamma_{S,\alpha}) \cdot R < 0$, then $(K_S + \Gamma_{S,\lambda}) \cdot R < 0$ and there is a rational curve C generating R such that

$$-2 \dim S \leq (K_S + \Gamma_{S,\lambda}) \cdot C < 0$$

If $\alpha \gg \lambda$ and if $M_S \cdot C > 0$, then $(K_S + \Gamma_{S,\alpha}) \cdot C > 0$, a contradiction. The same argument applies in each step of the LMMP because the Cartier index of M_S is preserved by the LMMP (by Cartier index of M_S we mean the smallest natural number n such that nM_S is Cartier). Therefore, the LMMP on $K_S + \Gamma_{S,\alpha}$ is also an LMMP on $K_S + \Gamma_{S,\alpha'}$ for any $\alpha' \geq \alpha$. In particular, if $\overline{S}$ is the good minimal model of $K_S + \Gamma_{S,\alpha}$ obtained by the LMMP above, then $\overline{S}$ is also a good minimal model of $K_S + \Gamma_{S,\alpha'}$ for any $\alpha' \geq \alpha$.

Step 4. Let $\overline{S}$ be the model constructed in Step 3. By construction, there is a commutative diagram



where μ is a resolution, e is a contraction, and c, d are also resolutions. By Step 3, we have $e^*d^*M_{\overline{S}} = e^*c^*M_S = M_W$ because M_S is numerically trivial on each step of the LMMP that produced $S \dashrightarrow \overline{S}$. Moreover, $D_{\overline{S}} + \alpha M_{\overline{S}}$ is semi-ample for any $\alpha \gg 0$.

Step 5. In general, $D_{\overline{S}}$ may not be nef. However, the LMMP on $K_S + \Gamma_{S,\alpha}$ in Step 3 is a partial LMMP on D_S , that is, each step of the LMMP on $K_S + \Gamma_{S,\alpha}$ is also a step of an LMMP on D_S but it may not be a full LMMP on D_S . In any case, there is a $\mathbb{Q}$ -divisor $\overline{P} \geq 0$ on $\overline{W}$ which is exceptional/ $\overline{S}$ and

$$c^*D_S = \overline{P} + d^*D_{\overline{S}}$$

Now put $G = e^*d^*D_{\overline{S}}$ and $\phi := \tau\mu$. Then, for some integer $l > 0$, we have natural isomorphisms

$$R(Y, lD_Y) \simeq R(S, lD_S) \simeq R(\overline{S}, lD_{\overline{S}}) \simeq R(W, lG)$$

By construction, $\alpha M_W + G$ is semi-ample for any $\alpha \gg 0$. Since $M_W \sim_{\mathbb{Q}} 0/Y$, we deduce that G is semi-ample over Y . Moreover,

$$\begin{aligned} \phi^*D_Y + \mu^*R^- &\sim_{\mathbb{Q}} \mu^*h^*(K_S + \Delta_S + L_S + bM_S) + \mu^*R^+ \\ &\sim_{\mathbb{Q}} e^*c^*D_S + \mu^*R^+ \\ &= e^*\overline{P} + G + \mu^*R^+ \end{aligned}$$

Put $D'_Y := \phi_*(e^*\overline{P} + G + \mu^*R^+)$. Since $e^*\overline{P} + G + \mu^*R^+ \geq 0$, we get $D'_Y \geq 0$, and since R^- is exceptional/ Y , we get $D'_Y \sim_{\mathbb{Q}} D_Y$. So,

$$\phi^*D'_Y + \mu^*R^- \sim_{\mathbb{Q}} e^*\overline{P} + G + \mu^*R^+$$

and by the negativity lemma we have equality

$$\phi^*D'_Y + \mu^*R^- = e^*\overline{P} + G + \mu^*R^+$$

Let $N = \phi^*D'_Y - G$. Then,

$$\phi_*N = \phi_*(N + \mu^*R^-) = \phi_*(e^*\overline{P} + \mu^*R^+) \geq 0$$

On the other hand, we know that G is nef over Y so N is anti-nef over Y . Therefore, by the negativity lemma we deduce $N \geq 0$ hence $\phi^*D'_Y \geq G$. Now replace D_Y with D'_Y and this completes the proof of the proposition. $\square$

Proof. (of Theorem 1.2) This follows from Proposition 3.1 using the same notation. $\square$

4. FUJINO-GONGYO CONJECTURE

As mentioned earlier the smoothness assumption of f in Theorem 1.3 cannot be removed. If $f : X \rightarrow Z$ is a surjective morphism between smooth projective varieties with $-K_X$ nef, then Zhang [Zh96] proved that either Z is uniruled or $\kappa(Z) = 0$. In particular, if $\dim Z = 1$, then $-K_Z$ is semi-ample. Therefore, a counter-example to Fujino-Gongyo's conjecture without the smoothness assumption on f is possible only when $\dim Z \geq 2$. We give few counter-examples of different flavours.

Example 4.1 Let B' be the $(4, 4)$ -divisor on $Z' = \mathbb{P}^1 \times \mathbb{P}^1$ which consists of 4 lines vertical with respect to the first projection and 4 lines vertical with respect to the second projection. Let $\sigma : Z \rightarrow Z'$ be the blow up of the 16 intersection points in B' . Let B be the strict transform of B' . Then $\sigma^*(B') = B + \sum_{i=1}^{16} 2E_i$, where E_i are the irreducible exceptional divisors over the 16 points. So B is an even divisor in Z , i.e., $B \sim 2L$ for some divisor L in Z . Let $f : X \rightarrow Z$ be the double cover ramified over B . We will show that $-K_X$ is semi-ample but $-K_Z$ is not semi-ample.

By construction, $K_{Z'}^2 = 8$ and from $K_Z = \sigma^*K_{Z'} + \sum_{i=1}^{16} E_i$ we get $K_Z^2 = -8$. Therefore, $-K_Z$ is not nef, and thus not semi-ample. Since $f : X \rightarrow Z$ is the double cover ramified over B , $K_X \sim_{\mathbb{Q}} f^*(K_Z + L)$. On the other hand,

$$K_Z + L \sim_{\mathbb{Q}} \sigma^*K_{Z'} + \frac{1}{2}(\sigma^*(B') - B) + \frac{1}{2}B = \sigma^*(K_{Z'} + \frac{1}{2}B') \sim_{\mathbb{Q}} 0$$

So $-K_X \sim_{\mathbb{Q}} 0$ is semi-ample. In this example f is a flat morphism but not a contraction.

Example 4.2 Let r and s be positive integers. Let $\mathcal{E} = \mathcal{O}_{\mathbb{P}^s} \oplus \mathcal{O}_{\mathbb{P}^s}(1)^{r+1}$ and let $Z_{r,s}$ be the smooth $(r + s + 1)$ -dimensional variety $\mathbb{P}(\mathcal{E})$. The projection $\pi_{r,s} : Z_{r,s} \rightarrow \mathbb{P}^s$ has a section σ with image $P_{r,s} \subset Z_{r,s}$ corresponding to the trivial quotient $\mathcal{E} \rightarrow \mathcal{O}_{\mathbb{P}^s}$. Let $f : X_{r,s} \rightarrow Z_{r,s}$ be the blow-up of $P_{r,s}$. One can check by calculating intersection numbers and using Kleiman's ampleness criterion that when $r > s$ the divisor $-K_{X_{r,s}}$ is ample but $-K_{Z_{r,s}}$ is not nef. For details see [De01] Example 1.36 and Example 3.16 (2). In this example f is a contraction but not a flat morphism.

For the convenience of the reader we reproduce another counter-example due to Hiroshi Sato given in [FG11], Example 4.6. In this case f would be both a contraction and a flat morphism.

Example 4.3 (H. Sato) Let Σ be the fan in $\mathbb{R}^3$ whose rays are generated by

$$\begin{aligned} x_1 &= (1, 0, 1), x_2 = (0, 1, 0), x_3 = (-1, 3, 0), x_4 = (0, -1, 0), \\ y_1 &= (0, 0, 1), y_2 = (0, 0, -1), \end{aligned}$$

and their maximal cones are

$$\begin{aligned} &\langle x_1, x_2, y_1 \rangle, \langle x_1, x_2, y_2 \rangle, \langle x_2, x_3, y_1 \rangle, \langle x_2, x_3, y_2 \rangle, \\ &\langle x_3, x_4, y_1 \rangle, \langle x_3, x_4, y_2 \rangle, \langle x_4, x_1, y_1 \rangle, \langle x_4, x_1, y_2 \rangle. \end{aligned}$$

Let Δ be the fan obtained from Σ by successive star subdivisions along the rays spanned by

$$z_1 = x_2 + y_1 = (0, 1, 1)$$

and

$$z_2 = x_2 + z_1 = 2x_2 + y_1 = (0, 2, 1).$$

We can see that $V = X_\Sigma$, the toric threefold corresponding to the fan Σ with respect to the lattice $\mathbb{Z}^3 \subset \mathbb{R}^3$, is a $\mathbb{P}^1$ -bundle over $Z = \mathbb{P}^1(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(3))$. The $\mathbb{P}^1$ -bundle structure $V \rightarrow Z$ is induced by the projection $\mathbb{Z}^3 \rightarrow \mathbb{Z}^2 : (x, y, z) \mapsto (x, y)$. The toric variety $X = X_\Delta$ corresponding to the fan Δ was obtained by successive blow-ups from V . The maximal cones of X_Δ are:

$$\begin{aligned} \tau_1 &= \langle x_1, x_2, z_2 \rangle, \tau_2 = \langle x_1, z_1, z_2 \rangle, \tau_3 = \langle x_1, z_1, y_1 \rangle, \\ \tau_4 &= \langle x_3, y_1, z_1 \rangle, \tau_5 = \langle x_3, z_2, z_1 \rangle, \tau_6 = \langle x_3, z_2, x_2 \rangle, \\ \sigma_3 &= \langle x_1, x_2, y_2 \rangle, \sigma_4 = \langle x_2, x_3, y_2 \rangle, \sigma_5 = \langle x_3, x_4, y_1 \rangle, \\ \sigma_6 &= \langle x_3, x_4, y_2 \rangle, \sigma_7 = \langle x_4, x_1, y_1 \rangle, \sigma_8 = \langle x_4, x_1, y_2 \rangle. \end{aligned}$$

To check $-K_X$ is semi-ample, we can either apply the base-point-free criterion on toric varieties (see [CLS] Theorem 6.1.7), or the fact that on toric varieties, semi-ampleness of a divisor is equivalent to nefness (see [CLS] Theorem 6.3.12). Here we apply the base-point-free criterion (see Remark 4.4).

We have $-K_X = \sum_{i=1}^4 D_{x_i} + D_{y_1} + D_{y_2} + D_{z_1} + D_{z_2}$, where D_u denotes the invariant Cartier divisor corresponding to the ray generated by the point u . The Cartier data associated to $-K_X$ are as follows:

$$\begin{aligned} m_{\tau_1} &= m_{\tau_6} = m_{\sigma_3} = m_{\sigma_4} = (-2, -1, 1), & m_{\tau_2} &= m_{\tau_3} = (0, 0, -1), \\ m_{\tau_4} &= m_{\tau_5} = (1, 0, -1), & m_{\sigma_5} &= (4, 1, -1), & m_{\sigma_6} &= (4, 1, 1), \\ m_{\sigma_7} &= (0, 1, -1), & m_{\sigma_8} &= (-2, 1, 1). \end{aligned}$$

One can check that all m_{τ_i}, m_{σ_j} are in P_{-K_X} . So $-K_X$ is base point free. Moreover, $-K_X$ is big as X is a projective toric variety. So X is a toric weak Fano manifold. The morphism $f : X \rightarrow Z$ induced by the projection $\mathbb{Z}^3 \rightarrow \mathbb{Z}^2$ is a flat morphism onto Z since every fiber of f is one-dimensional.

On the other hand, $-K_Z$ is not nef since $-K_Z \cdot C < 0$ where $C \subset Z$ is the image of the section of $Z \rightarrow \mathbb{P}^1$ corresponding to the trivial quotient $\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(3) \rightarrow \mathcal{O}_{\mathbb{P}^1}$.

Remark 4.4 Here we recall some basic facts about toric varieties.

(1) Base point freeness on toric varieties: We follow the notation of [CLS]. Let X_Σ be a complete toric variety of dimension n and let $D = \sum_\rho a_\rho D_\rho$ be a torus-invariant Cartier divisor on X_Σ . We define a polytope

$$P_D = \{m \in M_{\mathbb{R}} \mid \langle m, u_\rho \rangle \geq -a_\rho \text{ for all } \rho \in \Sigma(1)\} \subset M_{\mathbb{R}}$$

Let $\{m_\sigma\}_{\sigma \in \Sigma(n)}$ be the Cartier data associated to D , where $m_\sigma \in M$ with $\langle m_\sigma, u_\rho \rangle = -a_\rho$, for all $\rho \in \sigma(1)$. Then, D is base point free if and only if $m_\sigma \in P_D$ for all $\sigma \in \Sigma(n)$.

(2) Kodaira dimension: Now assume that D is nef. Then, the Kodaira dimension $\kappa(D) = \dim P_D$. In particular, if X_Σ is projective and $D = -K_{X_\Sigma}$ is nef, then it is not difficult to see that $\dim P_D = n$ hence D is nef and big.

Proof. (of Theorem 1.3) By Lemma 2.4 of [FG11], we can assume that f is a contraction: this is achieved by taking the Stein factorisation which preserves the smoothness assumption. We may assume that $\dim X > \dim Z$ otherwise the theorem holds trivially. Pick a closed point $z \in Z$. Since $-K_X$ is semi-ample, for some $m > 1$ we can find a divisor $D \in |-mK_X|$ such that if we put $B = \frac{1}{m}D$ then we have: $K_X + B \sim_{\mathbb{Q}} 0$, (X, B) is klt, and $D = \text{Supp } B$ is smooth (note that it is also possible to have $B = 0$ depending on the situation). Since f is smooth, we can choose D so that there is a neighbourhood U of z such that $D = \text{Supp } B$ is relatively smooth over U . In particular, this ensures that the discriminant part $B_Z = 0$ near z .

Let $\sigma: Y \rightarrow Z$ be an Ambro model for f and (X, B) . Then, possibly after shrinking U around z , we have $M_Y = \sigma^*M_Z$ over U by [FG11], Remark 4.3, or by [Ko07], Proposition 8.4.9 (3) and Theorem 8.5.1. Since $K_X + B \sim_{\mathbb{Q}} 0$,

$$K_Z + B_Z + M_Z \sim_{\mathbb{Q}} 0 \text{ and } K_Y + B_Y + M_Y \sim_{\mathbb{Q}} 0$$

and

$$K_Y + \Delta_Y + M_Y \sim_{\mathbb{Q}} D_Y := E$$

where Δ_Y and E are as in the Introduction. Since E is exceptional/ Z , D_Y is the only effective $\mathbb{Q}$ -divisor in its $\mathbb{Q}$ -linear equivalence class. Moreover, we have $\kappa(K_Y + \Delta_Y/T) \geq 0$ since

$$\kappa(K_Y + \Delta_Y + M_Y) = \kappa(D_Y) \geq 0$$

where $Y \rightarrow T$ is again as in the Introduction.

By Theorem 1.2, there is a resolution $\phi: W \rightarrow Y$ and a $\mathbb{Q}$ -divisor $G \geq 0$ such that $\alpha M_W + G$ is semi-ample for any $\alpha \gg 0$ and $G \leq \phi^*D_Y = \phi^*E$. In particular, this means that $\alpha M_W + G$ is semi-ample over U for $\alpha \gg 0$. Since over U we have $M_W = \phi^*\sigma^*M_Z$, we deduce that G is semi-ample over U . But G is effective and exceptional over U hence by the negativity lemma it is zero over U , that is, G is mapped into $Z \setminus U$. Thus, we can find a rational number $\alpha \gg 0$ and a $\mathbb{Q}$ -divisor $P_W \geq 0$ such that $\alpha M_W + G \sim_{\mathbb{Q}} P_W$ and that P_W does not contain any irreducible component of $(\sigma\phi)^{-1}\{z\}$. But P_W restricted to $(\sigma\phi)^{-1}\{z\}$ is numerically trivial so P_W does not intersect $(\sigma\phi)^{-1}\{z\}$. Therefore,

$$\alpha M_Z \sim_{\mathbb{Q}} P_Z := \sigma_*\phi_*P_W \geq 0$$

and z does not belong to $\text{Supp } P_Z$ which implies that the stable base locus of M_Z does not contain z . By construction, $\text{Supp } B_Z$ also does not contain z . Therefore, z is not in the stable base locus of $-K_Z$. Since z was chosen arbitrarily, we deduce that $-K_Z$ is semi-ample. $\square$

REFERENCES

- [Am99] F. Ambro; *The Adjunction Conjecture and its applications*. arXiv:math/9903060v3.
- [Am04] F. Ambro; *Shokurov's boundary property*, J. Differential Geom. 67 (2004), no. 2, 229–255.
- [Am05] F. Ambro; *The moduli b -divisor of an lc-trivial fibration*. Compos. Math. 141 (2005), no. 2, 385–403.
- [BCHM] C. Birkar, P. Cascini, C. Hacon, J. McKernan; *Existence of minimal models for varieties of log general type*. J. Amer. Math. Soc. 23 (2010), 405–468.
- [CLS] D. Cox, J. Little, H. Schenck; *Toric varieties*, Graduate Studies in Mathematics, vol 124.
- [De01] Olivier Debarre; *Higher-Dimensional Algebraic Geometry*, Universitext, Springer-Verlag, New York, 2001.
- [F03] O. Fujino; *A canonical bundle formula for certain algebraic fiber spaces and its applications*. Nagoya Math. J. Vol. 172 (2003), 129–171.
- [FG12] O. Fujino, Y. Gongyo; *On images of weak Fano manifolds II*. arXiv:1201.1130v1.
- [FG11] O. Fujino, Y. Gongyo; *On images of weak Fano manifolds*. Math. Zeit. 270 (2011), 531–544.
- [FM00] O. Fujino, S. Mori; *A canonical bundle formula*. J. Differential Geometry 56 (2000), no. 1, 167–188.
- [Ka91] Y. Kawamata; *On the length of an extremal rational curve*, Invent. Math. 105 (1991), 609–611.
- [Ka97] Y. Kawamata; *Subadjunction of log canonical divisors for a variety of codimension 2*, Contemporary Mathematics 207 (1997), 79–88.
- [Ka98] Y. Kawamata; *Subadjunction of log canonical divisors, II*, Amer. J. Math. 120 (1998), 893–899.
- [Ko07] J. Kollár; *Kodaira's canonical bundle formula and adjunction*. In "Flips for 3-folds and 4-folds", Oxford University Press (2007).
- [KMM92] J. Kollr, Y. Miyaoka, S. Mori; *Rational connectedness and boundedness of Fano manifolds*. J. Differential Geom. 36 (1992), no 3, 765–779.
- [KM98] J. Kollár, S. Mori; *Birational Geometry of Algebraic Varieties*. Cambridge University Press (1998).
- [LTZZ10] S.S.Y Lu, Y. Tu, Q. Zhang, Q. Zheng; *On semistability of Albanese maps*. Manuscripta Math., 131 (2010) no. 3-4, pp. 531–535.
- [M93] Y. Miyaoka; *Relative deformations of morphisms and applications to fibre spaces*. Comment. Math. Univ. St. Paul. 42 (1993), no 1, 1-7.
- [PSh09] Yu. G. Prokhorov, V. V. Shokurov; *Towards the second main theorem on complements*. J. Algebraic Geom. 18 (2009), 151–199.
- [Zh96] Qi. Zhang; *On projective manifolds with nef anticanonical bundles*. J. Reine Angew. Math. 478 (1996), 57–60.

DPMMS, Centre for Mathematical Sciences,
 Cambridge University,
 Wilberforce Road,
 Cambridge, CB3 0WB,
 UK
 email: c.birkar@dpmms.cam.ac.uk

Institute of Mathematics,
Chinese Academy of Sciences,
No. 55 Zhonguancun East Road,
Haidian District, Beijing, 100190,
China
email: yifeichen@amss.ac.cn